SLIDE INTO WINTER SAVINGS

Our winter offers with our flexible voucher option, our all-in-one package and flexible service. At StarSEQ we know that every project is unique.

1-Step 16S and ITS Microbiome analysis promotion and voucher option

<u>1-Step Bacterial microbiome analysis or 1-Step Fungal microbiome analysis:</u>



- 16S V4 region, Primer combination: 515F 806R¹
 16S V4-V5 region, Primer combination: 515F 909R¹
- **16S** V3-V4 region, Primer combination: $341F 806R^4$
- **ITS** ITS 1 region, Primer combination: ITS1F ITS2²



Price example:

Package A: 288	samples only	26,52 € per sample and one region
Package B: 192	samples only	31,54 € per sample and one region
Package C: 96	samples only	47,08 € per sample and one region
Package D: 48	samples only	76,17 € per sample and one region



Price examples, all other sample sizes are possible. We charge per sample. The price is a combination of a fixed price for the run and a variable price for the libraries depending on the number of samples.

No shared runs, customer samples are sequenced exclusively on one MiSeq run!

Eukaryotes (microbial) analysis:

V8-V9 region, Primer combination: 18S-1422f (5'-ataacaggtctgtgatgccct-3') - 18S_1797R (5'-ccttcygcaggttcacctac-3')⁵

V9 region, Primer combination: Illumina_Euk_1391f (5'-GTACACACCGCCCGTC-3') – Illumina_EukBr_1510r (5'-TGATCCTTCTGCAGGTTCACCTAC-3'), blocking primer optional³;

V9 region samples could not be combined with ITS, V8-V9 or 16S samples and need to be sequenced in a seperate MiSeq run because of smaller amplicon size!



Send us your gDNA samples and make use of our "All in One" service:

- Quality control
- Single step amplicon generation with reduced bias
- Double indexing, quality check, quantification, normalization and pooling of amplicons
- Illumina MiSeq / NextSeq 2000[™] sequencing package A-D: 2 x 300 nt paired-end sequencing with V3 chemistry
- Output package A-D: 20-30 M reads (including 25 % PhiX to balance the composition of bases)
- De-multiplexing of reads
- Data delivery via FTP server



^{1,2,3,4,5} Primer sequence references: Please see page 4!



SLIDE INTO WINTER SAVINGS

Our winter offers with our flexible voucher option, our all-in-one package and flexible service. At StarSEQ we know that every project is unique.

2-Step 16S and ITS Microbiome analysis

2-Step Bacterial and Archaea Microbiome analysis:

16S, V1-V2 region, Primer combination 27F (5'-AGAGTTTGATCMTGGCTCAG-3') – 338R (5'-TGCTGCCTCCCGTAGGAGT-3')

2-Step Fungal (18S) microbiome analysis:

ITS, ITS 1/2 region, Primer combination: ITS-u2 (5'-GAAYCATCGARTCTTTGAACGC-3') – ITS-p4 (5'-CCGCTTAKTGATATGCTTAAA-3')

ITS, ITS 2 region, Primer combination: ITS3F (5'-GCATCGATGAAGAACGCAGC-3') – ITS4R (5'-TCCTCCGCTTATTGATATGC-3')



2-Step Barcoding and Custom Amplicon analysis

2-Step Barcoding:

cytochrome b, cytb1 (5'-ccatccaacatctcagcatgatgaaa-3') – cytb2 (5' -gcccctcagaatgatatttgtcctca-3')

cytochrome oxidase I, CO I E (5'-ccagagattagagggaatcagtg-3') – CO 1 F (5'cctgcaggaggaggagagcc-3')

or 2-Step Custom Amplicons:

design of custom primers with Illumina specific overhangs

Request for offer: microbiome@starseq.com







Optional: gold standard bioinformatic analysis of 16S/18S/ITS sequences (QIIME2) The bioinformatics analysis pipeline consists of quality control, pre-processing of reads, taxonomy identification and visualization, calculation of alpha and beta-diversity metrics.

Request for offer: microbiome@starseq.com







Prices without tax. Cannot be combined with other promotions or discounts. Valid until 31.03.2025



Our winter offers with our flexible voucher option, our all-in-one
package and flexible service. At StarSEQ we know that every project is unique.

16S and ITS Microbiome analysis



Sequencing of your ready to load 16S/ITS/18S libraries

- Illumina MiSeq / NextSeq 2000[™] sequencing: 2 x 300 nt paired-end sequencing with V3 chemistry
- Output 20-30 M reads (including 25 % PhiX to balance the composition of bases)
- De-multiplexing of reads

- Free 16S metagenomics analysis (Illumina App)
- Data delivery via FTP server

Only 3300 € for MiSeq sequencing





Optimize your research with our 16S/ITS workflow solutions

From sample prep to library prep to sequencing and bioinformatics. StarSEQ offers a complete and comprehensive workflow for various 16S/ITS applications. Sequencing of the hypervariable regions of the 16S rRNA gene is the gold-standard for identification and classification of bacterial communities within a given sample.



StarSEQ is using established gold-standard primer sequences to target regions covering the hypervariable regions V3-V4, V4 and V4-V5. In contrast to common used methods, StarSEQ has designed its own single-step technique for simultaneous amplification, indexing and adaptoring of samples to reduce the bias caused during PCR.







Prices without tax. Cannot be combined with other promotions or discounts. Valid until 31.03.2025.

SLIDE INTO WINTER SAVINGS

Our winter offers with our flexible voucher option, our all-in-one
package and flexible service. At StarSEQ we know that every project is unique.

🔪 16S and ITS Microbiome analysis

Applications:

- Environmental Metagenomics (soil, water, air, biofilms, complex organic communities)
- Human or Animal Microbiome (skin, stool, gut, blood, swab)
- Monitoring of animal health (e.g. alternatively or supplementarily to FELASA-Test)
- Sterility Monitoring
- Detection of Contamination
- Monitoring of Biogas Plant
- Biosafety Monitoring
- Food Quality
- Clinical Samples . . .



You have no time/staff for routine preparation of gDNA from your collected samples?

As an additional service we offer also the purification of gDNA starting from $18 \in$.

Primer sequence references:

¹ Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology*, *18*(5), 1403–1414. https://doi.org/10.1111/1462-2920.13023

¹ Walters, W., Hyde, E. R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., ... Knight, R. (2016). Improved Bacterial 165 rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems, 1*(1), e00009-15. https://doi.org/10.1128/mSystems.00009-15

² White, T. J., T. Bruns, S. Lee, and J. W. Taylor. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pp. 315-322 In: PCR Protocols: A Guide to Methods and Applications. Academic Press, New York, NY. Gardes, M., and T. D. Bruns. 1993. ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. Mol. Ecol. 2: 113-118.

³ Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W., & Huse, S. M. (2009). A Method for Studying Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable Regions of Small-Subunit Ribosomal RNA Genes. *PLOS ONE, 4*(7), e6372. Retrieved from https://doi.org/10.1371/journal.pone.0006372

⁴ Klindworth A., Pruesse E., Schweer T., Peplies J., Quast C., Horn M., et al. . (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and nextgeneration sequencing-based diversity studies. Nucleic Acids Res. 41, 1–11. 10.1093/nar/gks808 see also https://support.illumina.com/downloads/16s_metagenomic_sequencing_library_preparation.html

⁵ Bradley, I. M., Pinto, A. J., & Guest, J. S. (2016). Design and Evaluation of Illumina MiSeq-Compatible, 185 rRNA Gene-Specific Primers for Improved Characterization of Mixed Phototrophic Communities. Applied and Environmental Microbiology, 82(19), 5878 LP-5891. https://doi.org/10.1128/AEM.01630-16

see also https://support.illumina.com/downloads/16s_metagenomic_sequencing_library_preparation.html







¹ Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb Ecol 75:129–137. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. Environ Microbiol. 2016;18: 1403-1414.